

Letter to the Editor: Will We All Become Electrosensitive?

ÖRJAN HALLBERG¹ AND GERD OBERFELD²

¹Hallberg Independent Research, Trångsund, Sweden

²Public Health Department Salzburg, Salzburg, Austria

Dear Editor,

Each year an increasing number of people claim to suffer from electrosensitivity (see, e.g., compilation of references given in Table 1), also known as being electrically hypersensitive (EHS). There are also other diseases, such as fibromyalgia and burn-out syndrome, that have symptoms similar to those exhibited by people suffering from electrosensitivity.

In Sweden, electrosensitivity is recognized as a handicap, but there is still controversy surrounding the diagnosis of the disease. The mainstream view by governmental and medical authorities is that this handicap is a psychological phenomenon with no basis in physical or medical mechanisms (Swedish National Board of Health and Welfare, SNBHW, 1995), whereby perpetuating the misconception that only a small fraction of the population is concerned about electrosensitivity or the proximity of new radio transmission masts.

The number of reported cases of electrosensitivity has been steadily increasing since it was first documented in 1991. Data presented here are estimates and are based on large sample inquiries where different sets of questions have been used. To determine whether the statistics indicate a sub-population of electrosensitivity or if the total population is at stake, we plotted reported prevalence estimates over time in a normal distribution diagram (Table 1 and Figure 1).

Contrary to the views of mainstream medical authorities, Figure 1 shows that the group of electrosensitive people around the world, including Sweden, is not just a small fraction that deviates from the rest of the healthy population. Instead, it points at the possibility that electrosensitivity will be more widespread in the near future. The extrapolated trend indicates that 50% of the population can be expected to become electrosensitive by the year 2017.

Data presented here were collected in Austria, Germany, Great Britain, Ireland, Sweden, Switzerland, and the United States.

Address correspondence to Örjan Hallberg, Hallberg Independent Research, Polkavägen 14B, Trångsund 142 65, Sweden; E-mail: oerjan.hallberg@swipnet.se

Table 1
Estimated prevalence of electrosensitive people in different years and countries

Measured year	% EI sensitive	Country, reported year	Ref. No.
1985	0.06	Sweden 1991 (0.025–0.125%)	National Encyclopedia Sw., 1991
1994	0.63	Sweden 1995	Anonymous est., 1994
1995	1.50	Austria 1995	Leitgeb N. et al., 1995, 2005
1996	1.50	Sweden 1998	SNBHW, Env. report, 1998
1997	2.00	Austria 1998	Leitgeb N. et al., 1998, 2005
1997	1.50	Sweden 1999	Hillert L. et al., 2002
1998	3.20	California 2002	Levallois P., 2002
1999	3.10	Sweden 2001	SNBHW, Env. report, 2001
2000	3.20	Sweden 2003	Sw Labour Union Sif, 2003
2001	6.00	Germany 2002	Schroeder E., 2002
2002	13.30	Austria 2003 (7.6–19%)	Spiß B., 2003
2003	8.00	Germany 2003	Infas, 2003
2003	9.00	Sweden 2004	Elöverkänsligas Riksförbund, 2005
2003	5.00	Schweiz 2005	Bern, Medicine Social, 2005
2003	5.00	Ireland 2005	This is London, 2005
2004	11.00	England 2004	Fox E., 2004
2004	9.00	Germany 2005	Infas, 2004
2017	50.00	Extrapolated to 50%	




Figure 1. The prevalence (%) of people around the world who consider themselves to be electrosensitive, plotted over time in a normal distribution graph. The endpoint at 50% is an extrapolated value. Variation explained is 91%, the endpoint not included.

References

- Anonymous estimate: 50,000 in 1994.
- Elöverkänsligas Riksförbund. (2005). *Funktionshindret Elöverkänslighet*.
- Fox, E. (2004). *Rapporteurs Report*. WHO Workshop on Electrical Hypersensitivity, Oct. 25–27.
- Hillert, L., Berglind, N., Arnetz, B. B., Bellander, T. (2002). *Scand J Work Environ Health*. 28(1):33–41.
- Infas. (2003). Ermittlungen der Befürchtungen und Ängste der breiten Öffentlichkeit hinsichtlich möglicher Gefahren der hochfrequenten elektromagnetischen Felder des Mobilfunks—jährliche Umfragen. B. f. Strahlenschutz. Bonn, Institut für angewandte Sozialwissenschaft GmbH: 1–34.
- Infas – Institut für angewandte Sozialwissenschaft GmbH Friedrich-Wilhelm-Straße 18, 53113 Bonn. Ermittlung der Befürchtungen und Ängste der breiten Öffentlichkeit hinsichtlich möglicher Gefahren der hochfrequenten elektromagnetischen Felder des Mobilfunks—jährliche Umfragen—Abschlussbericht über die Befragung im Jahr 2004.
- Leitgeb, N. (1995). Electrosensibilität. *VBÖ J* 1:51–55.
- Leitgeb, N. (1998). Electromagnetic hypersensitivity. *Proc Int Workshop on EMF and Non-Specific Health Symp.* pp. 8–16, Graz, Austria.
- Leitgeb, N., Schröttner, J., Böhm, M. (2005). Does “electromagnetic pollution” cause illness? An inquiry among Austrian general practitioners. *Wien Med Wochenschr*, 153:237–241.
- Levallois, P. et al. (2002). Study of self-reported hypersensitivity to electromagnetic fields in California. *Environ Health Persp* 110(4):619–623.
- Medicine Social et Preventive de l’Unirvesité de Berne, 2005.
- Proof of mobile health risk (2005). *This is London*, Feb. 9.
- Schroeder, E. (2002). Stakeholder-Perspektiven zur Novellierung der 26.BlmSchV. Ergebnisse der bundesweiten Telefonumfrage im Auftrag des Bundesamtes für Strahlenschutz (BfS).
- Spiß, B. (2003). Pilotstudie zu Mobilfunkstrahlung und Gesundheit—Modellierung der Immission mit den Programmen NIRView und CORLA. *Diplomarbeit an der Naturwissenschaftlichen Fakultät der Universität Salzburg*, Oktober 2003, Salzburg, Austria.
- Swedish Labour Union Sif Estimate, 2003.
- Swedish National Board of Health and Welfare. (1995). *Elektriska och Magnetiska fält och Hälsoeffekter. Report, 1*.
- Swedish National Board of Health and Welfare. (1998). *Environmental Health Report*.
- Swedish National Board of Health and Welfare. (2001). *Environmental Health Report*.
- Swedish National Encyclopedia. (1991). *Nationalencyklopedin. elöverkänslighet*, 5.